

UC Irvine

UC Irvine Previously Published Works

Title

Draft genome sequences of gammaproteobacterial methanotrophs isolated from lake washington sediment.

Permalink

<https://escholarship.org/uc/item/1s11m560>

Journal

Genome announcements, 3(2)

ISSN

2169-8287

Authors

Kalyuzhnaya, Marina G
Lamb, Andrew E
McTaggart, Tami L
et al.

Publication Date

2015-03-01

DOI

10.1128/genomea.00103-15

Peer reviewed

Draft Genome Sequences of Gammaproteobacterial Methanotrophs Isolated from Lake Washington Sediment

Marina G. Kalyuzhnaya,^{a*} Andrew E. Lamb,^{b*} Tami L. McTaggart,^{b*} Igor Y. Oshkin,^{b*} Nicole Shapiro,^c Tanja Woyke,^c Ludmila Chistoserdova^b

Department of Microbiology^a and Department of Chemical Engineering,^b University of Washington, Seattle, Washington, USA; DOE Joint Genome Institute, Walnut Creek, California, USA^c

* Present address: Marina G. Kalyuzhnaya, Biology Department, San Diego State University, San Diego, California, USA; Andrew E. Lamb, Northeastern University, Boston, Massachusetts, USA; Tami L. McTaggart, University of California Irvine, Irvine, California, USA; Igor Y. Oshkin, Winogradsky Institute of Microbiology, Russian Academy of Sciences, Moscow, Russia.

The genomes of *Methylosarcina lacus* LW14^T (=ATCC BAA-1047^T = JCM 13284^T), *Methylobacter* sp. strain 21/22, *Methylobacter* sp. strain 31/32, *Methylomonas* sp. strain LW13, *Methylomonas* sp. strain MK1, and *Methylomonas* sp. strain 11b were sequenced and are reported here. All the strains are obligately methanotrophic bacteria isolated from the sediment of Lake Washington.

Received 27 January 2015 Accepted 2 February 2015 Published 12 March 2015

Citation Kalyuzhnaya MG, Lamb AE, McTaggart TL, Oshkin IY, Shapiro N, Woyke T, Chistoserdova L. 2015. Draft genome sequences of gammaproteobacterial methanotrophs isolated from Lake Washington sediment. *Genome Announc* 3(2):e00103-15. doi:10.1128/genomeA.00103-15.

Copyright © 2015 Kalyuzhnaya et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marina G. Kalyuzhnaya, mkalyuzhnaya@mail.sdsu.edu.

Lake Washington sediment has served for decades as a model system for environmental studies on aerobic methane oxidation (1–8). A great deal of information has been accumulated regarding methane consumption rates (1, 3) and the composition of microbial communities that consume methane (4–8). A number of methanotroph species have been isolated in pure culture (3). Here, we announce the draft genome sequences of six obligately methanotrophic bacteria belonging to the family *Methylococcaceae* (Table 1).

Strains *Methylomonas* sp. LW13 and *Methylosarcina lacus* LW14^T were isolated in 1999 (3), and strain LW14^T has been formally described (9). Strains *Methylomonas* sp. 11b, *Methylomonas* sp. MK1, *Methylobacter* sp. 21/22, and *Methylobacter* sp. 31/32 were isolated from a sample collected in 2011 (7, 8). The *Methylobacter* strains are psychrophilic and do not grow at temperatures of >24°C, while the remaining strains grow well at 30°C (3, 9). DNA preparations were obtained using the phenol-chloroform

method (9). The draft genome sequences were generated at the Department of Energy (DOE) Joint Genome Institute (JGI), Walnut Creek, CA, USA, using the Illumina platform (10) or Pacific Biosciences (PacBio) technology (Table 1). The raw reads were assembled using HGAP (version 2.1.1; PacBio data) (11) or All-Paths, version 39750 (12) and/or Velvet, version 1.1.05 (Illumina data). All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov>. Genome annotation was performed using Prodigal (13), followed by a round of manual curation using GenePRIMP (14) for the genomes in <20 scaffolds. The predicted coding sequences were translated and used to search the National Center for Biotechnology Information (NCBI) nonredundant database and the UniProt, TIGRFam, Pfam, KEGG, COG, and InterPro databases. Additional gene prediction analysis was performed within the Integrated Microbial Genomes (IMG) platform (15). The genome statistics are shown in Table 1.

TABLE 1 Strains described, accession numbers, and general genome statistics

Strain	Sample collection date	NCBI accession no.	Sequencing platform (genome coverage [×])	No. of scaffolds (no. of contigs)	G+C content (%)	Size (Mb)	Gene count ^a	pMMO/sMMO ^b	Nif ^c	Hyd ^d
<i>Methylobacter</i> sp. 21/22	2011	JMLA000000000	PacBio (79)	4 (4)	49.5	4.7	4,239	2/0	+	+
<i>Methylobacter</i> sp. 31/32	2011	JPOH000000000	PacBio (104)	2 (2)	49.2	5.0	4,669	2/0	+	+
<i>Methylomonas</i> sp. 11b	2011	AZXX000000000	Illumina (755)	1 (2)	51.4	5.4	5,086	3/1	+	+
<i>Methylomonas</i> sp. MK1	2011	AQOV000000000	PacBio/Illumina (79)	5 (5)	51.5	5.2	4,851	3/0	+	+
<i>Methylomonas</i> sp. LW13	1999	JNLB000000000	Illumina (300)	42 (42)	51.8	5.2	4,806	2/1	+	+
<i>Methylosarcina lacus</i> LW14 ^T	1999	AZUN000000000	Illumina (912)	1 (1)	54.7	4.4	4,047	1/0	—	—

^a Number of genomic objects (coding sequences [CDSs], fragmented CDSs [fCDS], rRNA, tRNA, miscellaneous RNA [miscRNA]).

^b Number of gene clusters encoding particulate methane monooxygenase (pMMO) or soluble methane monooxygenase (sMMO).

^c Nitrogenase gene cluster. +, present; —, absent.

^d Hydrogenase (Nif) gene cluster.

As typical obligate methanotrophs, all the strains encode particulate methane monooxygenase for methane oxidation. In addition, soluble methane monooxygenase gene clusters were identified in the genomes of *Methylomonas* sp. strain 11b and *Methylomonas* sp. strain LW13. The complete gene inventories for the ribulose monophosphate pathway (both KDPG [2-keto-3-deoxy-6-phosphogluconate] and FBA [fructose 1,6-bisphosphate aldolase] variants [16]) are present in all genomes. All three *Methylomonas* strains, as well as *M. lacus* LW14^T, also possess complete sets of genes for the serine cycle, while in the *Methylobacter* species, the genes coding for one enzyme, phosphoenolpyruvate carboxylase, are not identifiable. None of the strains encode ribulose-1,5-bisphosphate carboxylase/oxygenase or its homologues. With the exception of *M. lacus* LW14^T, all the organisms encode functions for nitrogen fixation and hydrogen production/utilization. Respiratory nitrate-nitrite reductases are encoded only in the *Methylobacter* sp. genomes, while nitrite/nitrous oxide reductases are encoded only in the *Methylomonas* sp. and *M. lacus* LW14^T genomes. With the availability of their genomic sequences, these diverse *Methylococcaceae* isolates present prospective models for studying methanotrophy in freshwater lake sediments.

Nucleotide sequence accession numbers. The genome sequences have been deposited in GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under award no. DE-SC0005154 and DE-SC-0010556, and by the National Science Foundation under award no. MCB-0950183.

The work conducted by the U.S. Department of Energy Joint Genome Institute was supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231.

REFERENCES

- Lidstrom ME, Somers L. 1984. Seasonal study of methane oxidation in Lake Washington. *Appl Environ Microbiol* 47:1255–1260.
- Kuivila KM, Murray JW, Devol AH, Lidstrom ME, Reimers CE. 1988. Methane cycling in the sediments of Lake Washington. *Limnol Oceanogr* 33:571–581. <http://dx.doi.org/10.4319/lo.1988.33.4.0571>.
- Auman AJ, Stolyar S, Costello AM, Lidstrom ME. 2000. Molecular characterization of methanotrophic isolates from freshwater lake sediment. *Appl Environ Microbiol* 66:5259–5266. <http://dx.doi.org/10.1128/AEM.66.12.5259-5266.2000>.
- Costello AM, Auman AJ, Macalady JL, Scow KM, Lidstrom ME. 2002. Estimation of methanotroph abundance in a freshwater lake sediment. *Environ Microbiol* 4:443–450. <http://dx.doi.org/10.1046/j.1462-2920.2002.00318.x>.
- Nercissian O, Noyes E, Kalyuzhnaya MG, Lidstrom ME, Chistoserdova L. 2005. Bacterial populations active in metabolism of C₁ compounds in the sediment of Lake Washington, a freshwater lake. *Appl Environ Microbiol* 71:6885–6899. <http://dx.doi.org/10.1128/AEM.71.11.6885-6899.2005>.
- Kalyuzhnaya MG, Lapidus A, Ivanova N, Copeland AC, McHardy AC, Szeto E, Salamov A, Grigoriev IV, Suciu D, Levine SR, Markowitz VM, Rigoutsos I, Tringe SG, Bruce DC, Richardson PM, Lidstrom ME, Chistoserdova L. 2008. High-resolution metagenomics targets specific functional types in complex microbial communities. *Nat Biotechnol* 26:1029–1034. <http://dx.doi.org/10.1038/nbt.1488>.
- Beck DA, Kalyuzhnaya MG, Malfatti S, Tringe SG, Glavina Del Rio T, Ivanova N, Lidstrom ME, Chistoserdova L. 2013. A metagenomic insight into freshwater methane-utilizing communities and evidence for cooperation between the *Methylococcaceae* and the *Methylophilaceae*. *PeerJ* 1:e23. <http://dx.doi.org/10.7717/peerj.23>.
- Oshkin IY, Beck DA, Lamb AE, Tchesnokova V, Benuska G, McTaggart TL, Kalyuzhnaya MG, Dedysh SN, Lidstrom ME, Chistoserdova L. 21 October 2014. Methane-fed microbial microcosms show differential community dynamics and pinpoint taxa involved in communal response. *ISME J*. <http://dx.doi.org/10.1038/ismej.2014.203>.
- Kalyuzhnaya MG, Stolyar SM, Auman AJ, Lara JC, Lidstrom ME, Chistoserdova L. 2005. *Methylosarcina lacus* sp. nov., a methanotroph from Lake Washington, Seattle, and emended description of the genus *Methylosarcina*. *Int J Syst Evol Microbiol* 55:2345–2350. <http://dx.doi.org/10.1099/ijs.0.63405-0>.
- Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
- Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U. S. A.* 108:1513–1518. <http://dx.doi.org/10.1073/pnas.1017351108>.
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11: <http://dx.doi.org/10.1186/1471-2105-11-119>.
- Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <http://dx.doi.org/10.1038/nmeth.1457>.
- Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T, Huntemann M, Anderson I, Billis K, Varghese N, Mavromatis K, Pati A, Ivanova NN, Kyrpides NC. 2014. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res* 42:D560–D567. <http://dx.doi.org/10.1093/nar/gkt963>.
- Anthony C. 1982. *The biochemistry of methylotrophs*. Academic Press, Zürich, Switzerland.